



Preliminary gravimetric characterization of the Nova Colinas impact structure, Parnaíba basin, Brazil: Implications for hydrocarbon accumulation

Pietro Demattê Avona*¹; Alvaro Penteado Crósta¹; Marcos Alberto Rodrigues Vasconcelos²; Fernando Lessa Pereira¹.

¹IG/UNICAMP, ²IGeo/UFBA

Copyright 2023, SBGf - Sociedade Brasileira de Geofísica.

This paper was prepared for presentation during the 18th International Congress of the Brazilian Geophysical Society held in Rio de Janeiro, Brazil, 16-19 October 2023.

Contents of this paper were reviewed by the Technical Committee of the 18th International Congress of the Brazilian Geophysical Society and do not necessarily represent any position of the SBGf, its officers or members. Electronic reproduction or storage of any part of this paper for commercial purposes without the written consent of the Brazilian Geophysical Society is prohibited.

Abstract

Nova Colinas is the newest confirmed complex impact structure in Brazil, making it the ninth of its type known in the country. It is located in the Parnaíba sedimentary basin, in SW Maranhão State. This nearly circular structure has its center at 7°9'33"S and 46°6'30"W and a diameter between 6 to 7 km, with an estimated age of approximately 200 Ma based on stratigraphic constraints. Shock deformation features found at NC include planar fractures and planar deformation features in quartz, as well as shatter cones in basalt. NC is one of the rare examples on the Earth's surface of a meteoritic impact structure formed in mixed volcanic-sedimentary rocks. As the volcanic rocks comprise basalt, the study of NC may provide by analogy valuable insights for characterizing planetary geological processes, once basalt is a predominant lithology in planetary bodies such as the Moon, Mars, among others. Geological phenomena related to crater formation processes, such as fracturing, brecciation and hydrothermal alteration, can promote the buildup of hydrocarbon deposits. Impact structures with diameters similar to NC like Calvin (6 km) and Red Wing Creek (9 km), both located in the USA, are well-documented impact structures with associated hydrocarbon accumulations. Geophysics play a fundamental role in identifying and interpreting the subsurface characteristics of large impact structures which are important for analyzing the potential for oil and gas accumulations. A comprehensive geophysical characterization of the NC impact structure is underway, including the analysis of magnetometric, seismic and gravity data, in addition to numerical simulations. Here, we discuss the results of a preliminary gravimetric characterization of NC. We compared Bouguer anomaly maps with different slab densities aiming to reduce the topographic effect on the collected data. A previous work shows that a density of 2,05 g/cm³ would fit better than the standard slab density of 2,67 g/cm³ for this region. The final Bouguer maps show similar results in terms of intensity and geometry of the anomalies. The results allowed a clear distinction between two different domains: (i) one in the NE area exhibiting a high gravity peak of -77.4 mGal, probably related to the basalt of the Mosquito Formation, and (ii) another in the SW area with lower gravity values reaching -86.5 mGal, likely related to the sandstones of the Sambaíba Formation. Although the structure's boundaries are not well defined by the Bouguer anomaly, it exhibits a nearly circular anomaly in the central area with a high gravity of approximately -78.8 mGal, likely related to a central structural uplift. Due to the impact-related (shock) effects on the target rocks, it is expected to exhibit a central low gravity signature typical of complex craters, which does not occur at NC. A reasonable explanation for this lack at the center is related to deeper densest rocks within the stratigraphic sequence which were uplifted due to the impact. These densest rocks comprise the Motuca and Pedra de Fogo Formations, composed in the area predominantly by intercalations between heterolytic/sandstones rocks and intercalations between siltstone/silicite with some anhydrite layers, respectively.