

A DECADE OF AIRGLOW OBSERVATION OF MESOSPHERIC FRONTS OVER THE BRAZILIAN EQUATORIAL REGION

Emanuel Barbosa de Carvalho ^{1*}, Igo Paulino ¹ and Cristiano Max Wrasse ²

¹Universidade Federal de Campina Grande - UFCG, Campina Grande, PB, Brazil

²Instituto Nacional de Pesquisas Espaciais - INPE, São José dos Campos, SP, Brazil

*Corresponding author: emanuelbarbosadecarvalho@gmail.com

ABSTRACT. Mesospheric fronts were observed over São João do Cariri (7.4°S, 36.5°W), in the northeast region of Brazil, from September 2011 to June 2021, using an all-sky airglow imager. The occurrence of the mesospheric fronts presented a well-defined seasonality with its in the Spring period of ten consecutive years. In addition, an anisotropy was observed in the preferential propagation direction, in which most of the front propagated to the northeast. These results suggest that the seasonality and anisotropy can be produced due to either the location of sources or the filtering process associated with the wind system. Other important aspects like the statistics of trailing waves and ripples observed simultaneously were reported as well.

Keywords: mesospheric bore, atmosphere dynamics, airglow imager, gravity waves, ripples.

INTRODUCTION

Mesospheric fronts are gravity waves with a well-defined front, pronounced in amplitude, spatially extended, followed or not by trailing waves and not necessary in a wave-like structure (Brown et al., 2004). They can be classified as bore, wall or simply front (Dewan and Picard, 1998). This phenomenon has been mostly observed using airglow cameras in low (e.g., Isler et al., 1997; Medeiros et al., 2001, 2005, 2016, 2018; Fechine et al., 2005, 2009; Narayanan et al., 2009; Carvalho et al., 2017), middle (e.g., She et al., 2004; Smith et al., 2005) and high (e.g., Nielsen et al., 2006; Bageston et al., 2011a,b; Giongo et al., 2018) latitudes. The mesospheric fronts can propagate for long horizontal distances into ducts with small dissipation of energy (Dewan and Picard, 2001); thus, they are important in the transport of energy and momentum for long distances in the atmosphere. The formation of these ducts depends on the background atmosphere conditions and thermodynamics. A thermal duct can be produced due to changes in the temperature profile structures (e.g., Sarkhel et al., 2022); Doppler ducts are produced due to the vertical wind shear (e.g., Fechine et al., 2009); and a dual duct can appear when both temperature and wind are relevant for the formation (e.g., Bageston et al., 2011a).

Most of the recent advances on the mesospheric fronts came from case studies using multi instrument observations, in which the focus is primarily to describe the events, the propagation conditions of the atmosphere and the formation of the ducts (Medeiros et al., 2016, 2018; Smith et al., 2017; Hozumi et al., 2018; Mondal et al., 2021; Thurairajah et al., 2021) and/or identify possible sources of them (Mehta et al., 2017; Giongo et al., 2018). Theoretical studies have also been developed giving some directions for further investigations (e.g., Dewan and Picard, 1998, 2001; Laughman et al., 2009). Another important topic of research on mesospheric fronts is the appearance of convective and dynamic instabilities in the Mesosphere and Lower Thermosphere (MLT) when they interact with the background atmosphere (e.g., Fritts et al., 2020, and references therein).

In contrast, publications using long-term observations are rare; consequently, some aspects of mesospheric front events are poorly understood, such as seasonality and solar cycle dependency. Based on it, the present work aims to give some relevant statistical findings on mesospheric fronts using airglow observations over the northeast of Brazil from 2011 to 2021. For instance, the seasonality of the occurrence, preferential propagation direction, and presence of trailing waves and ripples simultaneously to the front will

be reported and discussed. INSTRUMENTATION AND OBSERVATION aims to describe the methodology and define some important aspects used in this study. In RESULTS AND DISCUSSION, the results are presented with the characteristics of the registered fronts, preferential propagation direction and seasonal occurrence of the phenomena. In SUMMARY AND CONCLUSION, the main contributions of this work are highlighted and the results are summarized.

INSTRUMENTATION AND OBSERVATION

In this work, near infrared OH airglow images were collected at São João do Cariri ($7.4^{\circ}S, 36.5^{\circ}W$) from 23 September 2011 to 15 June 2021 from a Charge-Coupled Device (CCD) all-sky imager. Images with an integration time of 15 seconds and a temporal sample of about 2 minutes were used. The imager uses a fisheye lens which allows us to make images with 180° of field of view. The instrument is equipped with a telecentric system of lenses which projects the light orthogonal to the interference filters mounted on a filter wheel. After that, the lights pass through a system of lenses to reconstruct the image on the CCD chip. The CCD temperature is set to cooled by a Peltier cooler at $-70^{\circ}C$. The whole system is controlled by a microcomputer. The CCD has 6.45 cm^2 of area with resolution of 2048×2048 pixels, back-illuminated and binned on-chip down to a resolution of 512×512 pixels to enhance the signal to noise ratio. It has high quantum efficiency ($\approx 70\%$ in the visible), high linearity (0.05%), low thermal (0.5 electrons/pixel/second) and reading (15 electron/pixel) noises. Details about the imager have been published elsewhere (e.g., [Paulino et al., 2011](#)).

Using the whole period of observations (23 September 2011 - 15 June 2021), characteristics of the mesospheric fronts were tabulated, such as: (i) the start and end time of the observations, (ii) the period of cloudiness, (iii) the interval of time when the fronts crossed the field of view of the imager. Double checking of the observed fronts was done using OI6300 that comes from different altitudinal ranges, it was considered a mesospheric front if the structure was observed only in the OH images. Additionally, in the confirmed fronts, it was investigated the presence of (a) trailing waves following the fronts; (b) the presence of ripples simultaneously to the passage of the front in the images and (c) the uniformity of the background of the images. The fronts were identified using the conditions stated by [Brown et al. \(2004\)](#).

Figure 1 shows two examples of mesospheric fronts. Figure 1a shows an example of a front followed by trailing waves 28 June 2019 over São João do Cariri, and Figure 1b shows an example with the presence of ripples while a front was propagating on 16 September 2015.

The next step of the methodology was the calculation of the amount of Valid Observation Hours (VOH)

per season, the seasonal average, and their respective standard deviations. VOH is defined as the amount of hours in which the sky was cloudless. The usage of this quantity as stated above aims to reduce possible misleads that could come out from the difference in the amount of time when it was actually possible to observe the sky for each season, since there is no uniformity between the seasons in this specific aspect.

The identification of the propagation directions was done by tracing a line perpendicularly to the front in the best observed image. Then, it was calculated the clockwise angle from the north (azimuth). Table 1 shows the conventions used.

Table 1: Conventions relative to the angles and directions of propagation of the bores.

Direction	Angle
North	Bigger than 337.5 and smaller than 22.5
Northeast	Between 22.5 and 67.5
East	Between 67.5 and 112.5
Southeast	Between 112.5 and 157.5
South	Between 157.5 and 202.5
Southwest	Between 202.5 and 247.5
West	Between 247.5 and 292.5
Northwest	Between 292.5 and 337.5

As this study was carried out in Brazil, the calendar data is based on Southern Hemisphere seasons. To determine the beginning and end of the seasons from 2011 to 2020, it was considered the convention provided by the Astronomy Department of the Institute of Astronomy, Geophysics and Atmospheric Sciences of Universidade de São Paulo - USP and, for the year of 2021, it was taken a date based on the other year dates of the beginning of the seasons.

RESULTS AND DISCUSSION

From 23 September 2011 to 15 June 2021, 246 fronts were observed. Regarding the season, as shown in Figure 2a, (i) 13.00% of the fronts occurred in Summer; (ii) 20.74% in Autumn; (iii) 30.08% in Winter; and (iv) 36.18% in Spring. Figure 2b shows that, out of the total amount of fronts, 87.8% of the fronts were followed by trailing waves, 19.9% of them occurred with the presence of ripples in the field of view of the imager and only 4.9% of the fronts had non uniform color when propagated through the images (a front was considered non uniform in its aspect when, for example, the front appeared in the images firstly as a bright band well pronounced in amplitude but as time passed this band became dark; the same kind of change may also occur when a dark band becomes a bright band as time passes.)




Figure 1: (a) Front propagating to North followed by trailing waves, registered on 28 June 2019; (b) Front propagating to East followed by trailing waves and with the presence of ripples in the all-sky field of view, registered on 16 September 2015 over São João do Cariri.

The fact that this amount of fronts are followed by trailing waves points out that the sources of these fronts must be far from the observatory, since the presence of trailing waves indicates that there is loss of energy.

The occurrence of ripples implies that the front could be interacting with the background producing dynamical and convective instabilities (e.g., Fritts and Rastogi, 1985; Medeiros et al., 2018). Given that in our data this phenomenon has low rates of occurrence, it is possible to point out a low frequency of these instabilities in this region associated with the passage of the mesospheric fronts.

Figure 3 shows the propagation direction of the fronts. For every season, at least 58.82% of the fronts were propagating to the North, Northeast, or Southeast. All-year data shows that fronts propagating to those directions represent 63.82% (16.67% to North, 26.42% to Northeast and 20.73% to Southeast) of the identified fronts. Therefore, it is possible to state Northeast, Southeast and North as the preferential propagation directions of mesospheric fronts over São João do Cariri. The fact appointed by this graphic can be related either to the location of the sources of the fronts or to the needed physical conditions for its canalization. Small and medium-scale gravity waves observed over São João do Cariri also presented a quite similar anisotropy in the propagation direction (Campos et al., 2016; Essien et al., 2018). Those authors also suggested the location of the sources and the filtering by the wind system as possible mechanisms to explain the observed anisotropy.

Looking at the seasonal average of occurrence by VOH in Figure 4, it is seen a considerable dispersion of the data, but with a clear growth over the seasons. The seasonal average of occurrence by VOH grows

and reaches its maximum value in Spring, which shows that the phenomena are more likely to occur in Spring than in other seasons. The seasonality of the occurrence of the fronts is also in agreement with other gravity waves observed at the same site (e.g., Essien et al., 2018), suggesting again that the sources of the fronts must be related to the source of other gravity wave structures.

SUMMARY AND CONCLUSION

Long-term observations of OH airglow images collected between September 2011 and June 2021 in the Northeast of Brazil revealed interesting characteristics of mesospheric fronts that are summarized as follows:

- The amount of fronts followed by trailing waves suggests sources far from the field of view of the imager while the low rates of ripple occurrence suggest that just a few fronts dissipate over São João do Cariri, once the ripple presence is related to the dissipation of mesospheric fronts;
- The identified preferential propagation directions were Northeast, Southeast and North. Fact that can be explained either by the location of the sources of these waves or by the conditions of canalization of these waves (or both). More studies are needed to determine the predominant factor that makes these directions preferential;
- The occurrence by VOH average grows over the seasons reaching its maximum average in Spring, pointing that there is a seasonality in the occurrence of the phenomena for the years 2011-2020.




Figure 2: General information about the mesospheric fronts.




Figure 3: Registered mesospheric fronts by propagation direction per season and all-year long.




Figure 4: Registered fronts per VOH by season. Each error bar corresponds to the standard deviation of ten values, each of them corresponding to the respective season of a year. The solid line represents the linear fit to the points.

ACKNOWLEDGEMENTS

The airglow images used in this study can be accessed online at the site of the Instituto Nacional de Pesquisas Espaciais - INPE (www2.inpe.br/climaespacial/portal/original-image-video). The table used to define the seasons of each year from 2011 to 2020 is available at www.iag.usp.br/astrografia/inicio-das-estacoes-do-ano. This work was developed with the financial support of the Conselho Nacional de Desenvolvimento Científico e Tecnológico - CNPq (grant number: 306063/2020-4), Fundação de Amparo à Pesquisa do Estado da Paraíba - PRONEX (Edital Universal) and Programa Institucional de Bolsas de Iniciação Científica, Universidade Federal de Campina Grande (PIBIC-UFCG/CNPq).

REFERENCES

- Bageston, J. V., C. M. Wrasse, P. P. Batista, R. E. Hibbins, D. Fritts, D. Gobbi, and V. F. Andrioli, 2011a, Observation of a mesospheric front in a thermal-doppler duct over King George Island, Antarctica: *Atmospheric Chemistry and Physics*, **11**, 12137–12147, doi: [10.5194/acp-11-12137-2011](https://doi.org/10.5194/acp-11-12137-2011).
- Bageston, J. V., C. M. Wrasse, R. E. Hibbins, P. P. Batista, D. Gobbi, H. Takahashi, V. F. Andrioli, J. Fechine, and C. M. Denardini, 2011b, Case study of a mesospheric wall event over Ferraz Station, Antarctica (62°S): *Annales Geophysicae*, **29**, 209–219, doi: [10.5194/angeo-29-209-2011](https://doi.org/10.5194/angeo-29-209-2011).
- Brown, L. B., A. J. Gerrard, J. W. Meriwether, and J. J. Makela, 2004, All-sky imaging observations of mesospheric fronts in OI 557.7 nm and broadband OH airglow emissions: analysis of frontal structure, atmospheric background conditions, and potential sourcing mechanisms: *Journal of Geophysical Research: Atmospheres*, **109**, D19104, doi: [10.1029/2003JD004223](https://doi.org/10.1029/2003JD004223).
- Campos, J. A. V., I. Paulino, C. M. Wrasse, A. F. de Medeiros, A. R. Paulino, and R. A. Buriti, 2016, Observations of small-scale gravity waves in the equatorial upper mesosphere: *Brazilian Journal of Geophysics*, **34**, 469–477, doi: [10.22564/rbgf.v34i4.876](https://doi.org/10.22564/rbgf.v34i4.876).
- Carvalho, A., I. Paulino, A. Medeiros, L. Lima, R. Buriti, A. Paulino, C. Wrasse, and H. Takahashi, 2017, Case study of convective instability observed in airglow images over the Northeast of Brazil: *Journal of Atmospheric and Solar-Terrestrial Physics*, **154**, 33–42, doi: [10.1016/j.jastp.2016.12.003](https://doi.org/10.1016/j.jastp.2016.12.003).
- Dewan, E. M., and R. H. Picard, 1998, Mesospheric bores: *Journal of Geophysical Research: Atmospheres*, **103**, 6295–6305, doi: [10.1029/97JD02498](https://doi.org/10.1029/97JD02498).
- Dewan, E. M., and R. H. Picard, 2001, On the origin of mesospheric bores: *Journal of Geophysical Research: Atmospheres*, **106**, 2921–2927, doi: [10.1029/2000JD900697](https://doi.org/10.1029/2000JD900697).
- Essien, P., I. Paulino, C. M. Wrasse, J. A. V. Campos, A. R. Paulino, A. F. Medeiros, R. A. Buriti, H. Takahashi, E. Agyei-Yeboah, and A. N. Lins, 2018, Seasonal characteristics of small- and medium-scale gravity waves in the mesosphere and lower thermosphere over the Brazilian equatorial region: *Annales Geophysicae*, **36**, 899–914, doi: [10.5194/angeo-36-899-2018](https://doi.org/10.5194/angeo-36-899-2018).
- Fechine, J., A. Medeiros, R. Buriti, H. Takahashi, and D. Gobbi, 2005, Mesospheric bore events in the equatorial middle atmosphere: *Journal of Atmospheric and Solar-Terrestrial Physics*, **67**, 1774–

- 1778, doi: [10.1016/j.jastp.2005.04.006](https://doi.org/10.1016/j.jastp.2005.04.006).
- Fechine, J., C. M. Wrasse, H. Takahashi, A. F. Medeiros, P. P. Batista, B. R. Clemesha, L. M. Lima, D. C. Fritts, B. Laughman, M. J. Taylor, P. D. Pautet, M. G. Mlynczak, and J. M. Russell, 2009, First observation of an undular mesospheric bore in a doppler duct: *Annales Geophysicae*, **27**, 1399–1406, doi: [10.5194/angeo-27-1399-2009](https://doi.org/10.5194/angeo-27-1399-2009).
- Fritts, D. C., N. Kaifler, B. Kaifler, C. Geach, C. Kjellstrand, B. P. Williams, S. D. Eckermann, A. J. Miller, M. Rapp, M. Limon, J. Reimuller, and L. Wang, 2020, Mesospheric Bore Evolution and Instability Dynamics Observed in PMC Turbo Imaging and Rayleigh Lidar Profiling Over Northeastern Canada on 13 July 2018: *Journal of Geophysical Research: Atmospheres*, **125**, e2019JD032037, doi: [10.1029/2019JD032037](https://doi.org/10.1029/2019JD032037).
- Fritts, D. C., and P. K. Rastogi, 1985, Convective and dynamical instabilities due to gravity wave motions in the lower and middle atmosphere: Theory and observations: *Radio Science*, **20**, 1247–1277, doi: [10.1029/RS020i006p01247](https://doi.org/10.1029/RS020i006p01247).
- Giongo, G. A., J. V. Bageston, P. P. Batista, C. M. Wrasse, G. D. Bittencourt, I. Paulino, N. M. Paes Leme, D. C. Fritts, D. Janches, W. Hocking, and N. J. Schuch, 2018, Mesospheric front observations by the OH airglow imager carried out at Ferraz Station on King George Island, Antarctic Peninsula, in 2011: *Annales Geophysicae*, **36**, 253–264, doi: [10.5194/angeo-36-253-2018](https://doi.org/10.5194/angeo-36-253-2018).
- Hozumi, Y., A. Saito, T. Sakanoi, A. Yamazaki, and K. Hosokawa, 2018, Undulating wave front of mesospheric bore: Space-borne observations by ISS-IMAP/VISI: *Atmospheric Chemistry and Physics Discussions*, **18**, 16399–16407, doi: [10.5194/acp-2018-383](https://doi.org/10.5194/acp-2018-383).
- Isler, J. R., M. J. Taylor, and D. C. Fritts, 1997, Observational evidence of wave ducting and evanescence in the mesosphere: *Journal of Geophysical Research: Atmospheres*, **102**, 26301–26313, doi: [10.1029/97JD01783](https://doi.org/10.1029/97JD01783).
- Laughman, B., D. C. Fritts, and J. Werne, 2009, Numerical simulation of bore generation and morphology in thermal and Doppler ducts: *Annales Geophysicae*, **27**, 511–523, doi: [10.5194/angeo-27-511-2009](https://doi.org/10.5194/angeo-27-511-2009).
- Medeiros, A., J. Fechine, R. Buriti, H. Takahashi, C. Wrasse, and D. Gobbi, 2005, Response of OH, O₂ and OI5577 airglow emissions to the mesospheric bore in the equatorial region of Brazil: *Advances in Space Research*, **35**, 1971–1975, doi: [10.1016/j.asr.2005.03.075](https://doi.org/10.1016/j.asr.2005.03.075).
- Medeiros, A., M. J. Taylor, H. Takahashi, P. Batista, and D. Gobbi, 2001, An unusual airglow wave event observed at Cachoeira Paulista (23°S): *Advances in Space Research*, **27**, 1749–1754, doi: [10.1016/S0273-1177\(01\)00317-9](https://doi.org/10.1016/S0273-1177(01)00317-9).
- Medeiros, A. F., I. Paulino, M. J. Taylor, J. Fechine, H. Takahashi, R. A. Buriti, L. M. Lima, and C. M. Wrasse, 2016, Twin mesospheric bores observed over Brazilian equatorial region: *Annales Geophysicae*, **34**, 91–96, doi: [10.5194/angeo-34-91-2016](https://doi.org/10.5194/angeo-34-91-2016).
- Medeiros, A. F., I. Paulino, C. M. Wrasse, J. Fechine, H. Takahashi, J. V. Bageston, A. R. Paulino, and R. A. Buriti, 2018, Case study of mesospheric front dissipation observed over the Northeast of Brazil: *Annales Geophysicae*, **36**, 311–319, doi: [10.5194/angeo-36-311-2018](https://doi.org/10.5194/angeo-36-311-2018).
- Mehta, D., A. J. Gerrard, Y. Ebihara, A. T. Weatherwax, and L. J. Lanzerotti, 2017, Short-period mesospheric gravity waves and their sources at the South Pole: *Atmospheric Chemistry and Physics*, **17**, 911–919, doi: [10.5194/acp-17-911-2017](https://doi.org/10.5194/acp-17-911-2017).
- Mondal, S., M. Sivakandan, S. Sarkhel, M. Sunil Krishna, M. G. Mlynczak, J. M. Russell, and G. Bharti, 2021, A case study of a thermally ducted undular mesospheric bore accompanied by ripples over the Western Himalayan region: *Advances in Space Research*, **68**, 1425–1440, doi: [10.1016/j.asr.2021.03.026](https://doi.org/10.1016/j.asr.2021.03.026).
- Narayanan, V. L., S. Gurubaran, and K. Emperumal, 2009, A case study of a mesospheric bore event observed with an all-sky airglow imager at Tirunelveli (8.7°N): *Journal of Geophysical Research: Atmospheres*, **114**, doi: [10.1029/2008JD010602](https://doi.org/10.1029/2008JD010602).
- Nielsen, K., M. J. Taylor, R. G. Stockwell, and M. J. Jarvis, 2006, An unusual mesospheric bore event observed at high latitudes over Antarctica: *Geophysical Research Letters*, **33**, L07803, doi: [10.1029/2005GL025649](https://doi.org/10.1029/2005GL025649).
- Paulino, I., H. Takahashi, A. Medeiros, C. Wrasse, R. Buriti, J. Sobral, and D. Gobbi, 2011, Mesospheric gravity waves and ionospheric plasma bubbles observed during the COPEX campaign: *Journal of Atmospheric and Solar-Terrestrial Physics*, **73**, 1575–1580, doi: [10.1016/j.jastp.2010.12.004](https://doi.org/10.1016/j.jastp.2010.12.004).
- Sarkhel, S., G. Stober, J. L. Chau, S. M. Smith, C. Jacobi, S. Mondal, M. G. Mlynczak, and J. M. Russell, 2022, A case study of a ducted gravity wave event over Northern Germany using simultaneous airglow imaging and wind-field observations: *Annales Geophysicae Discussions*, **40**, 179–190, doi: [10.5194/angeo-2021-48](https://doi.org/10.5194/angeo-2021-48).
- She, C. Y., T. Li, B. P. Williams, T. Yuan, and R. H. Picard, 2004, Concurrent OH imager and sodium temperature/wind lidar observation of a mesopause region undular bore event over Fort Collins/Platteville, Colorado: *Journal of Geophysical Research: Atmospheres*, **109**, D22107, doi: [10.1029/2004JD004742](https://doi.org/10.1029/2004JD004742).
- Smith, S. M., J. Friedman, S. Raizada, C. Tepley, J. Baumgardner, and M. Mendillo, 2005, Evidence of mesospheric bore formation from a breaking gravity wave event: simultaneous imaging and lidar measurements: *Journal of Atmospheric and Solar-Terrestrial Physics*, **67**, 345–356, doi: [10.1016/j.jastp.2004.11.008](https://doi.org/10.1016/j.jastp.2004.11.008).
- Smith, S. M., G. Stober, C. Jacobi, J. L. Chau,

M. Gerding, M. G. Mlynczak, and G. Umbriaco, 2017, Characterization of a double mesospheric bore over Europe: *Journal of Geophysical Research: Space Physics*, **122**, 9738–9750, doi: [10.1002/2017JA024225](https://doi.org/10.1002/2017JA024225).

Thurairajah, B., C. Y. Cullens, and S. M. Bailey, 2021, Characteristics of a mesospheric front observed in polar mesospheric cloud fields: *Journal of Atmospheric and Solar-Terrestrial Physics*, **218**, 105627, doi: [10.1016/j.jastp.2021.105627](https://doi.org/10.1016/j.jastp.2021.105627).

Carvalho, E.B.: analysis, Writing – original draft preparation; **Paulino, I.:** supervision, Writing – review & edition, analysis, interpretation; **Wrasse, C.M.:** experiment implementation, Writing – review & edition.

Received on December 31, 2021 / Accepted on November 30, 2022



Creative Commons attribution-type CC BY